REFERENCES

1. Moftat CA., Osseltion MD., Widdop B.,(2004), Clarke's Analysis of drugs and poisons , Pharmaceutical press, 3rd edition.

2. Wiley J., Connors AK., (1999), Pharmaceutical analysis 3rd edition,221-224

3.Jeffery GH., Basett J., Mendham J., Denny RC., (1996), Vogels Text book of Quantitatives chemical analysis 5th edition,383-393

4. Chitturi.S R.,Somannavar YS., Peruri BG., Nallapatis Sharma H.K., Budidest SR.,Handa VK.,Vurimindi HB.,(2011)., RP-HPLC methods for the determination of process impurities and degradation products of atazanavie sulphate drugs substance, journal of bio med. Analysis, 55(1), 31-47

5. Swetha Mallesh A., Ravindra reddy Y.,(2011), Method development of atazanavir and ritonavir in a combined dosage form by RP-HPLC method, International Journal of Pharmacy and technology, 3316-3334

6.KatarzynMichalik.,Zofiadrzazga.,Annamichnik.,Kaszuba.M.,(2007), Thermal stability study of the protease inhibitors, nelfinavir, mesylate and atazanavir sulphate, Journal of thermal analysis and calorimetry, 88(2), 401-404.

7. Janaki pathi P.,Ravendra reddy P., Appalaraju N.,(2011), Visible spectrophotometric estimation of atazanavir in pharmaceutical formulations , 4(2) 273-275

8.Estelle Cateau.,Nicolas Tournier.,Antoinedepuis.,Gwenael Le Moal.,Nicolas Venisse.,(2005), Abstract determination of atazanavir in human plasma using solid phase extraction and HPLC, Journal of pharmaceutical and bio medical analysis ,39 (3-4) , 791-795

9.Colombo S., Guijnard N.,Marzolinic.,Telenti A.,Biollaz J., Decosterd LA., (2004), determination of the new HIV-Protase inhibitor Atazanavir by Liquidchromatography atter solid phase extraction, Journal of chromatography B.analyst technol bio med life science, 810(1), 25-34

10.Jie Ling., Kin johnsen A.,Zhuang Miao.,Ashok rakhit.,Michel PantzeP.,MartaHomilton.,BertLumL.,Chandraprakash.,(2006),Metabolism and excretion of erlotinib , A small molecule inhibitor of epidermal growth factor receptor tyrosine kinase, in healthy male volunteers, The American society for pharmacology and experimental therapeutics,34(3),420-426

11.Vallur Rajesh.,Annupama B., Jagathi V.,Varaprasad K.,(2011), HPLC method for estimation of erlotinib hydro chloride as bulk drug, international journal of biological and medical research 2(1),433-435

12.Erin lepper R.,Sandra swain M.,Antoinette.,Jan R., Willian Figg D.,Alex sparreboom.,(2003),Liquid-Chromotographic determination of erlotinib, epidermal growth factor receptor tyrosine kinase inhibitor, journal of chromatography B, 796(1), 181-188

13.Pujeri S.S.,Khader A.M.A., Seetharamppa J.,(2009), validated stability indicating chromatographic method for the assay of erlotinib active pharmaceutical ingredient, enalitical letters 42(12),1867-1885

14Rajesh V.,Jagathi V.,sindhuri K.,Devalarao G., (2011), Spectro fluorimetric method for estimation of erlotinib hydrochloride in pure and pharmaceutical formulations, E.Journal of chemistry,8(S1),S304-S308

15.Chahbouni.,Den burger A Bsc., Bsc JCG.,Vos., Sinjewel RM., Wilhelnm ABsc., (2009), Simultaneous quantification of erlotinib, gefitinib and lmatinib in human plasma by liquid chromatography tandam mass spectro metric, Therapeutic drug monitoring , 31(6), 683-687

16.Padmalatha M., Syeda Kulsum.,Rahul C.,Timma reddy D.,Vidyasagar G., (2011)., Spectro photometric methods for the determination of erlotinib in pure and pharmaceutical dosage forms, international journal of pharmaceutical research and development ,3(6), 103-109

17.Mukhopadhyay S.,Rokade N.,Sawant L.,Nachane D.,Padita N.,(2011).,Validated RP-HPLC method for determination of related substances of prulifloxacin in tablet dosage form, research journal of pharmacy and technology , 4(9), 1379-1385

18.Deepak Porkhar.,Varsha Jadhav.,Sachin Gltolve.,Vilas rao kadam., (2010), Development and validation of spectro photometric method for estimation of prulifloxacin in tablet dosage form international journal of pharma technology and research, 2(1), 960-963

19.JunWen.,Zhenyu Zhu., Zhansing Hong.,Yiwen Wu., Yanfed.,Mei Lin.,Guorang Fan.,Yutian wu.,(2007) Determination of active metabolite of prulifloxacin in human plasma by HPLC with fluorescence detection , Chromatographia: an internation journal for rapid communication in chromatography, electro phorosis and associated techniques, 66(1), 37-41

20. Srinivasu K., Venkateshwara Rao J., Appala Raju N., Mukkanti., (2011), A validated RP HPLC method for the determination of Atazinavir in pharmaceutical dosage form, E journal of chemistry8(1), 453-456

21.Suddha sattya Dey., Vikram Reddy Y., Thirupathi reddy., Sudhir tsahoo., Murthy P. N., Subhasis Mohapata., Subhasis S.,(2010) method development and validation for the estimation of atazanavir in bulk and pharmaceutical dosage forms and its stress degradation studies using UV –VIS spectrophotometric method, International journal of pharma and biosciences 1(3), 1-12

22 .Vijayalakshmi A., Gowri Shankar D., Jeevani G., Samson Israel D., (2010) validated RP-HPLC method as a tool for the estimation of Prulifloxacin in pharmaceutical dosage form, international journal of pharma research and development 2(5)1-7.

23. Chaple D. R., Sambhare A. G., (2010) A validated stability indicating HPLC method for Prulifloxacin, International journal of pharmacy and Technology,1 (2),137-148

24. Lixia Guo., Meiling Qi., Xin Jin., Peng Wang., Huaiquing Zhao., (2006), determination of active metabolite of Prulifloxacin in human plasma by liquid chromatography- tandem mass spectrometry, journal of chromatography B, 832(2),280-285

25. Fouad chiadmi., Mathieu Duprez., Joel Schlatter., Jean-eudes Fontan., validation of a simple high performance liquid chromatography analysis method for determination of erlotinib in human plasma,European journal of hospital pharmacy,13(2),48-51
26. Kalyana Chakravarthy V., Gowri sankar D., (2011),development and validation of RPHPLC method for the estimation of erlotinib in bulk and its pharmaceutical formulation4(2),393-399
27. Dhanya B., Suganthi A., Sen A. K., Sahoo U., Seth A. K.,(2011), determination of doxazosin mesylate in tablets by RPHPLC, Indian journal of pharmaceutical sciences, 73(1), 120-122
28. Aydogmus , Zeynep, Barla, Asli(2009), spectrophotometric determination of doxazosin mesylaste in tablets by ion pair and charge-transfer complexation reactions, Jaoac Int92(1), 131-138
29. Qutab S. S., Razzaq S.N., Ashfaq M., Shuja Z. A., Khan I. U.,(2007) simple and sensitive LC UV method for simultaneous analysis of hydrochlorthiazide and Candesartan cilexetil in pharmaceutical formulations, ACTA Chromatographica, (19), 119-129
30. Subba Rao D.V., Radha krishanand P., Surya Narayana M.V., Himabindu V.,(2007),Stability indicating LC method for Candesartan celixetil,Chromatographia,66, 499-507
31. Claudius., malerczyk.,Bettina fuchs.,Gustav G.,Beiz.,Susanna Roll.,Volker herrman.,steffen magin G.,anette Hogemann.,Barbara voith.,Ernst Mutscbler.,(1998),Angiotensin Antagonism and plasma radio receptor kinetics of Candesartan in man,British Journal clinical pharmacol,45(6),567-573
32. Balamurli Krishna K., Syamasundar B., (2010), development and validation of high performance liquid chromatographic method for the simultaneous estimation of Candesartan cilexetil and hydrochlorthiazide in combined tablet dosage form, Der pharma chemical, 2(6), 231-237
33. Patel Jignesh,Dave J. B., Patel C.N., Patel dhrumil, (2010), Q –Analysis spectrophotometric methods for estimation of Candesartan cilexitil and hydrochlorthiazide in tablet dosage form, J. Chem. Pharma Res,2(3),10-14
34. Patil Banswaraj S., Rao raghavendra N. G., Jadhav Suvarna., Upendra K., Gada Mahesh M., (2011), estimation of cilexitil in bulk and tablet dosage form by UV spectrophotometric method, international journal of ayurdeva and pharmacy, 2(1),204-206
35. Chetan Detroja,.chavan s.,Krutika sawanth.,(2011),enhanced antihypertensive activity of candesartan cilexetil nanosuspspention, formulation characterization and pharmacodynamic study, scientia pharmaceutica sci pharm,79, 635-657
36. Sopan wadhe P., Dnyani Wadhe P., Bakal R.L., (2011),RPHPLC method for simultaneous determination of Drotaverine hydrochloride and paracetamol in tablet dosage form, International journal of Pharmaceutical research and development, 3(8), 145-150
37. Dahivelkar R.P.,Mahagan V.K., Bari S.B., Shirknedkar A.A., Fursule R. A., Surana S.J., (2007)simultaneous derivative and multi-component spectrophotometric determination of drotaverine hydrochloride and Mefenamic acid in tablets, Indian Journal of pharmaceutical sciences, 69(6), 812-814

38. Garg G., Swarnalatha Saraf, Saraf S., (2007), simultaneous estimation of mefenemic acid and ethamsylate in tablets, Indian Journal of pharmaceutical sciences, 69(2),279-28